ELSEVIER

Neuroscience Letters 322 (2002) 195–197

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 322

Aasly, J., see Farrer, M., 83
Adachi, Y., see Matsumoto, Y., 67
Adam, A., see Farrer, M., 83
Adams Jr., J.D., see Chang, M.L., 137
Aker, R.G., Özkara, Ç., Dervent, A., Yılmaz Onat, F., Enhancement of spike and wave discharges by microinjection of bicuculline into the reticular nucleus of rats with absence epilepsy, 71
Albin, R.L., see Lavedan, C., 57
Alderson, H.L., see Jenkins, T.A., 45
Amano, T., see Tomita, Y., 157
Angrilli, A., Penolazzi, B., Vespignani, F., De Vincenzi, M., Job, R., Ciccarelli, L., Palomba, D., Stegagno, L., Cortical brain responses to semantic incongruity and syntactic violation in Italian language: an event-related potential study, 5

Back, S.K., Sung, B., Hong, S.K., Na, H.S., A mouse model for peripheral neuropathy produced by a partial injury of the nerve supplying the tail, 153
Ballesteros, M.A., González, F., Morón, I., DeBrugada, I., Cándido, A., Gallo, M., Dissociation of the associative and visceral sensory components of taste aversion learning by tetrodotoxin inactivation of the parabrachial nucleus in rats, 169
Bartolini, A., see Galeotti, N., 145
Berg, M., see Farrer, M., 83
Bernstein, J.J., see DeGiorgio, L.A., 62
Biessmann, A., see Haas, J., 121
Bisceglio, G., see Farrer, M., 83
Bole, D.G., Hirata, K., Ueda, T., Prolonged depolarization of rat cerebral synaptosomes leads to an increase in vesicular glutamate content, 17
Brand, G., see Millot, J.-L., 79
Buchholtz, S., see Lavedan, C., 57
Butcher, S.P., see Macleod, M.R., 41

Cándido, A., see Ballesteros, M.A., 169
Cannizzaro, C., see Zeng, B.-Y., 33
Cavaletti, G., Pezzoni, G., Pisano, C., Oggioni, N., Sala, F., Zoia, C., Ferrarese, C., Marmiroli, P., Tredici, G., Cisplatin-induced peripheral neurotoxicity in rats reduces the circulating levels of nerve growth factor, 103
Cervero, F., see Galan, A., 37
Chan, P.H., see Chang, M.L., 137
Chang, M.L., Yang, J., Kem, S., Klaidman, L., Sugawara, T., Chan, P.H., Adams Jr., J.D., Nicotinamide and ketamine reduce infarct volume and DNA fragmentation in rats after brain ischemia and reperfusion, 137
Chen, J.-F., see Xu, K., 13
Choue, R.-W., see Jang, M.-H., 29
Chung, J.-H., see Jang, M.-H., 29
Ciccarelli, L., see Angrilli, A., 5
Compton, D., see Fidani, L., 192
Cookson, M.R., see O'Farrell, C., 9

Dass, B., see Zeng, B.-Y., 33
De Vincenzi, M., see Angrilli, A., 5
DeBrugada, I., see Ballesteros, M.A., 169
DeGiorgio, L.A., Manuelidis, L., Bernstein, J.J., Transient appearance of amyloid precursor protein plaques in the brain of thymectomized rats after human leptomeningeal cell grafts, 62
Dervent, A., see Aker, R.G., 71
Di Cesare Mannelli, L., see Galeotti, N., 145
Du, J.-L., Yang, X.-L., Bullfrog retinal bipolar cells may express heterogeneous glycine receptors at dendrites and axon terminals, 177

Elepfandt, P., Rupprecht, S., Schöning-Burkhardt, B., Volk, H.-D., Woiciechowsky, C., Oligodeoxynucleotides induce brain inflammation in rats when infused intracerebroventricularly, 107
Eller, M., Harro, J., Partial denervation of the locus coeruleus projections by treatment with the selective neurotoxin DSP-4 potentiates the long-term effect of parachloroamphetamine on 5-hydroxytryptamine metabolism in the rat, 53
Endo, T., Isa, T., Postsynaptic and presynaptic $GABA_B$ receptor-mediated inhibition in rat superficial superior colliculus neurons, 126

Farrer, M., Skipper, L., Berg, M., Bisceglio, G., Hanson, M., Hardy, J., Adam, A., Gwinn-Hardy, K., Aasly, J., The Tau H1 Haplotype is associated with Parkinson's disease in the Norwegian population, 83
Feng, Y.-P., Yang, K., Li, Y.-Q., Norepinephrine depresses the capsaicin-evoked miniature excitatory postsynaptic currents in substantia gelatinosa of the rat spinal cord, 99
Ferrarese, C., see Cavaletti, G., 103
Fidani, L., Compton, D., Hardy, J., Petersen, R.C., Tangalos, E., Mirtsou, V., Goulas, A., Wavrant De Vrieze, F., No association between the lipoprotein lipase S447X polymorphism and Alzheimer's disease, 192
Fujino, M., see Matsumoto, Y., 67
Fukuuchi, Y., see Tomita, Y., 157
Fukuyama, H., see Ino, T., 182
Furuya, K., see Kawahara, N., 91

Galan, A., Lopez-Garcia, J.A., Cervero, F., Laird, J.M.A., Activation of spinal extracellular signaling-regulated kinase-1 and -2 by intraplantar carrageenan in rodents, 37
Galeotti, N., Di Cesare Mannelli, L., Mazzanti, G., Bartolini, A., Ghelardini, C., Menthol: a natural analgesic compound, 145
Gallo, M., see Ballesteros, M.A., 169
Ghelardini, C., see Galeotti, N., 145
González, F., see Ballesteros, M.A., 169
Goulas, A., see Fidani, L., 192

PII: S0304-3940(02)00206-9

Gulley, J.M., Kosobud, A.E.K., Rebec, G.V., Amphetamine inhibits behavior-related neuronal responses in substantia nigra pars reticulata of rats working for sucrose reinforcement, 165
Gwinn-Hardy, K., see Farrer, M., 83

Haas, J., Storch-Hagenlocher, B., Biessmann, A., Wildemann, B., Inducible nitric oxide synthase and argininosuccinate synthetase: co-induction in brain tissue of patients with Alzheimer's dementia and following stimulation with β-amyloid 1–42 in vitro, 121
Han, H.C., see Hong, S.K., 21
Han, J.S., see Hong, S.K., 21
Hanson, M., see Farrer, M., 83
Hardy, J., see Farrer, M., 83
Hardy, J., see Fidani, L., 192
Harro, J., see Eller, M., 53
Hauss-Wegrzyniak, B., Wenk, G.L., Beta-amyloid deposition in the brains of rats chronically infused with thiorphan or lipopolysaccharide: the role of ascorbic acid in the vehicle, 75
Hiramatsu, T., see Xiong, G., 173
Hirata, K., see Bole, D.G., 17
Hirose, S., see Ino, T., 182
Honda, K., Koga, K., Moriyama, T., Koguchi, M., Takano, Y., Kamiya, H.-o., Intrathecal α_2 adrenoceptor agonist clonidine inhibits mechanical transmission in mouse spinal cord via activation of muscarinic M_1 receptors, 161
Hong, S.K., Han, J.S., Min, S.S., Hwang, J.M., Kim, Y.I., Na, H.S., Yoon, Y.W., Han, H.C., Local neurokinin-1 receptor in the knee joint contributes to the induction, but not maintenance, of arthritic pain in the rat, 21
Hong, S.K., see Back, S.K., 153
Hwang, J.M., see Hong, S.K., 21

Ino, T., Inoue, Y., Kage, M., Hirose, S., Kimura, T., Fukuyama, H., Mental navigation in humans is processed in the anterior bank of the parieto-occipital sulcus, 182
Inoue, Y., see Ino, T., 182
Isa, T., see Endo, T., 126
Itoh, T., see Matsumoto, Y., 67
Ivanova, E.A., see Popova, N.K., 1

Jang, M.-H., Shin, M.-C., Lim, B.-V., Chung, J.-H., Kang, H.-S., Kang, S.-A., Choue, R.-W., Kim, E.-H., Kim, C.-J., Nicotine administration decreases nitric oxide synthase expression in the hypothalamus of food-deprived rats, 29
Jenkins, T.A., Latimer, M.P., Alderson, H.L., Winn, P., Determination of acetylcholine and dopamine content in thalamus and striatum after excitotoxic lesions of the pedunculopontine tegmental nucleus in rats, 45
Jenner, P., see Zeng, B.-Y., 33
Job, R., see Angrilli, A., 5

Kage, M., see Ino, T., 182
Kamiya, H.-o., see Honda, K., 161
Kang, H.-S., see Jang, M.-H., 29
Kang, S.-A., see Jang, M.-H., 29
Kawahara, N., Kawai, K., Toyoda, T., Nakatomi, H., Furuya, K., Kirino, T., Cardiac arrest cerebral ischemia model in mice failed to cause delayed neuronal death in the hippocampus, 91
Kawai, K., see Kawahara, N., 91
Kem, S., see Chang, M.L., 137
Kim, C.-J., see Jang, M.-H., 29
Kim, E.-H., see Jang, M.-H., 29
Kim, Y.I., see Hong, S.K., 21
Kimura, T., see Ino, T., 182
Kirino, T., see Kawahara, N., 91
Kirov, R., Moyanova, S., Distinct sleep-wake stages in rats depend differentially on age, 134
Kiviniemi, K., see Vihla, M., 141
Klaidman, L., see Chang, M.L., 137
Kobari, M., see Tomita, Y., 157
Koga, K., see Honda, K., 161
Koguchi, M., see Honda, K., 161
Kõks, S., Planken, A., Luuk, H., Vasar, E., Cat odour exposure increases the expression of wolframin gene in the amygdaloid area of rat, 116
Kosobud, A.E.K., see Gulley, J.M., 165
Kurokawa, T., see Matsumoto, Y., 67

Laird, J.M.A., see Galan, A., 37
Latimer, M.P., see Jenkins, T.A., 45
Lavedan, C., Buchholtz, S., Nussbaum, R.L., Albin, R.L., Polymeropoulos, M.H., A mutation in the human neurofilament M gene in Parkinson's disease that suggests a role for the cytoskeleton in neuronal degeneration, 57
Li, Y.-Q., see Feng, Y.-P., 99
Lim, B.-V., see Jang, M.-H., 29
Lopez-Garcia, J.A., see Galan, A., 37
Luuk, H., see Kõks, S., 116

Macleod, M.R., Butcher, S.P., Nitric-oxide-synthase-mediated cyclic guanosine monophosphate production in neonatal rat cerebellar prisms is resistant to calcineurin inhibition, 41
Manuelidis, L., see DeGiorgio, L.A., 62
Marmiroli, P., see Cavaletti, G., 103
Matsumoto, Y., Watanabe, T., Adachi, Y., Itoh, T., Ohtaki, T., Onda, H., Kurokawa, T., Nishimura, O., Fujino, M., Galanin-like peptide stimulates food intake in the rat, 67
Matsu-ura, T., Nakadai, T., Oda, Y., Nagasu, T., Mikoshiba, K., Tamura, T.-a., Seizure-mediated accumulation of the β subunit of Ca^{2+}/calmodulin-dependent protein kinase II in nuclei of mouse brain cells, 149
Mazzanti, G., see Galeotti, N., 145
McGowan, E., see O'Farrell, C., 9
Menendez de la Prida, L., Suarez, F., Pozo, M.A., The effect of different morphological sampling criteria on the fraction of bursting cells recorded in the rat subiculum in vitro, 49
Mikoshiba, K., see Matsu-ura, T., 149
Millot, J.-L., Brand, G., Morand, N., Effects of ambient odors on reaction time in humans, 79
Milton, N.G.N., The amyloid-β peptide binds to cyclin B1 and increases human cyclin-dependent kinase-1 activity, 131
Min, S.S., see Hong, S.K., 21
Mirtsou, V., see Fidani, L., 192
Morand, N., see Millot, J.-L., 79
Moriyama, T., see Honda, K., 161
Morón, I., see Ballesteros, M.A., 169
Moyanova, S., see Kirov, R., 134
Mutoh, S., see Ochi, T., 95

Na, H.S., see Back, S.K., 153
Na, H.S., see Hong, S.K., 21
Nagao, S., see Xiong, G., 173
Nagasu, T., see Matsu-ura, T., 149
Nakadai, T., see Matsu-ura, T., 149
Nakatomi, H., see Kawahara, N., 91
Nishimura, O., see Matsumoto, Y., 67

Nussbaum, R.L., see Lavedan, C., 57

Ochi, T., Ohkubo, Y., Mutoh, S., Blockade of the antinociceptive effect of spinally administered kyotorphin by naltrindole in mice, 95
Oda, Y., see Matsu-ura, T., 149
O'Farrell, C., Pickford, F., Vink, L., McGowan, E., Cookson, M.R., Sequence conservation between mouse and human synphilin-1, 9
Ogawa, H., Wang, X.-D., Neurons in the cortical taste area receive nociceptive inputs from the whole body as well as the oral cavity in the rat, 87
Oggioni, N., see Cavaletti, G., 103
Ohkubo, Y., see Ochi, T., 95
Ohtaki, T., see Matsumoto, Y., 67
Ohtomo, M., see Tomita, Y., 157
Onda, H., see Matsumoto, Y., 67
Owen, A., see Zeng, B.-Y., 33
Özkara, Ç., see Aker, R.G., 71

Palomba, D., see Angrilli, A., 5
Pan, Y.-P., see Xu, X.-H., 25
Penolazzi, B., see Angrilli, A., 5
Petersen, R.C., see Fidani, L., 192
Pezzoni, G., see Cavaletti, G., 103
Pickford, F., see O'Farrell, C., 9
Pisano, C., see Cavaletti, G., 103
Planken, A., see Kõks, S., 116
Polymeropoulos, M.H., see Lavedan, C., 57
Popova, N.K., Ivanova, E.A., 5-HT_{1A} receptor antagonist p-MPPI attenuates acute ethanol effects in mice and rats, 1
Pozo, M.A., see Menendez de la Prida, L., 49

Rebec, G.V., see Gulley, J.M., 165
Rose, S., see Zeng, B.-Y., 33
Rupprecht, S., see Elepfandt, P., 107

Sakamoto, H., Ukena, K., Tsutsui, K., Dendritic spine formation in response to progesterone synthesized de novo in the developing Purkinje cell in rats, 111
Sala, F., see Cavaletti, G., 103
Salmelin, R., see Vihla, M., 141
Schiszler, I., see Tomita, Y., 157
Schöning-Burkhardt, B., see Elepfandt, P., 107
Schwarzschild, M.A., see Xu, K., 13
Shin, M.-C., see Jang, M.-H., 29
Shum, A.Y.-C., see Wang, J.-Y., 187
Skipper, L., see Farrer, M., 83
Stegagno, L., see Angrilli, A., 5
Storch-Hagenlocher, B., see Haas, J., 121
Suarez, F., see Menendez de la Prida, L., 49
Sugawara, T., see Chang, M.L., 137
Sung, B., see Back, S.K., 153

Takano, Y., see Honda, K., 161
Takeda, H., see Tomita, Y., 157
Tamura, T.-a., see Matsu-ura, T., 149
Tanahashi, N., see Tomita, Y., 157
Tangalos, E., see Fidani, L., 192
Tel, B.C., see Zeng, B.-Y., 33
Tomita, M., see Tomita, Y., 157
Tomita, Y., Tomita, M., Schiszler, I., Amano, T., Tanahashi, N., Kobari, M., Takeda, H., Ohtomo, M., Fukuuchi, Y., Repetitive concentric wavering spread of oligemia/hyperemia in the sensorimotor cortex accompanying K^+-induced spreading depression in rats and cats, 157
Toyoda, T., see Kawahara, N., 91
Tredici, G., see Cavaletti, G., 103
Tsutsui, K., see Sakamoto, H., 111

Ueda, T., see Bole, D.G., 17
Ukena, K., see Sakamoto, H., 111

Vasar, E., see Kõks, S., 116
Vespignani, F., see Angrilli, A., 5
Vihla, M., Kiviniemi, K., Salmelin, R., Auditory cortical activation in Finnish and Swedish speaking Finns: a magnetoencephalographic study, 141
Vink, L., see O'Farrell, C., 9
Volk, H.-D., see Elepfandt, P., 107

Wang, J.-Y., see Wang, J.-Y., 187
Wang, J.-Y., Shum, A.Y.-C., Wang, J.-Y., Hypoxia/reoxygenation induces cell injury via different mechanisms in cultured rat cortical neurons and glial cells, 187
Wang, X.-D., see Ogawa, H., 87
Wang, X.-L., see Xu, X.-H., 25
Watanabe, T., see Matsumoto, Y., 67
Wavrant De Vrieze, F., see Fidani, L., 192
Wenk, G.L., see Hauss-Wegrzyniak, B., 75
Wildemann, B., see Haas, J., 121
Winn, P., see Jenkins, T.A., 45
Woiciechowsky, C., see Elepfandt, P., 107

Xiong, G., Hiramatsu, T., Nagao, S., Corticopontocerebellar pathway from the prearcuate region to hemispheric lobule VII of the cerebellum: an anterograde and retrograde tracing study in the monkey, 173
Xu, K., Xu, Y.-H., Chen, J.-F., Schwarzschild, M.A., Caffeine's neuroprotection against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine toxicity shows no tolerance to chronic caffeine administration in mice, 13
Xu, X.-H., Pan, Y.-P., Wang, X.-L., mRNA expression alterations of inward rectifier potassium channels in rat brain with cholinergic impairment, 25
Xu, Y.-H., see Xu, K., 13

Yang, J., see Chang, M.L., 137
Yang, K., see Feng, Y.-P., 99
Yang, X.-L., see Du, J.-L., 177
Yılmaz Onat, F., see Aker, R.G., 71
Yoon, Y.W., see Hong, S.K., 21

Zeng, B.-Y., Dass, B., Owen, A., Rose, S., Cannizzaro, C., Tel, B.C., Jenner, P., 6-Hydroxydopamine lesioning differentially affects α-synuclein mRNA expression in the nucleus accumbens, striatum and substantia nigra of adult rats, 33
Zoia, C., see Cavaletti, G., 103